



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/78, C12N 9/68, C07K 14/515, A61K 47/48	A1	(11) International Publication Number: WO 99/48923 (43) International Publication Date: 30 September 1999 (30.09.99)
(21) International Application Number: PCT/US99/06246 (22) International Filing Date: 22 March 1999 (22.03.99) (30) Priority Data: 09/046,985 24 March 1998 (24.03.98) US (71) Applicants: THE CHILDREN'S MEDICAL CENTER CORPORATION [US/US]; 300 Longwood Avenue, Boston, MA 02115 (US). YISSUM RESEARCH DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; Jabotinsky Street 46, 91042 Jerusalem (IL). (72) Inventor: BEN-SASSON, Shmuel, A.; Epstein Street 3, 96555 Jerusalem (IL). (74) Agents: WAGNER, Richard, W. et al.; Hamilton, Brook, Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA 02421 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: MULTIVALENT LIGANDS WHICH MODULATE ANGIOGENESIS <div style="text-align: center; margin: 20px 0;"> $\text{B} - \left(\text{L} - \text{P} \right)_n \quad (I)$ </div> (57) Abstract <p>Disclosed are novel multivalent ligands represented by structural formula (I) wherein B is a multilinker backbone, n is an integer from two to about twenty, each L is a covalent bond or linking group, each P is a peptide having from about 10 to about 30 amino acid residues. At least two of the peptides P are a peptide derivative of an AHR of an angiogenic protein, a hybrid peptide, a peptide derivative of a hybrid peptide or a combination thereof. Each peptide and each linker or covalent bond is independently chosen. The disclosed multivalent ligands can be used to modulate angiogenesis in a mammal. Also disclosed are novel peptide derivatives of an AHR of an angiogenic protein, novel hybrid peptides, peptide derivatives of the novel hybrid peptides and polypeptide multivalent ligands thereof.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

MULTIVALENT LIGANDS WHICH MODULATE ANGIOGENESIS

BACKGROUND OF THE INVENTION

Anti-angiogenic therapies are potential treatments for a wide variety of diseases, including cancer, arteriosclerosis, obesity, arthritis, duodenal ulcers,
5 cardiovascular disorders and abnormal ocular neovascularization caused, for example, by diabetes (Folkman, *Nature Medicine* 1:27 (1995) and Folkman, Seminars in Medicine of the Beth Israel Hospital, Boston, *New England Journal of Medicine*, 333:1757 (1995)). Anti-angiogenic therapies are thought to act by inhibiting the formation of new blood vessels.

10 Pro-angiogenic therapies are potential treatments for promoting wound healing and for stimulating the growth of new blood vessels to by-pass occluded ones. Thus, pro-angiogenic could potentially augment or replace by-pass surgeries and balloon angioplasty (PTCA).

The full potential of anti-angiogenic and pro-angiogenic therapies, together
15 referred to as "angiogenic therapies", has yet to be fully realized. One reason is because of the shortage of agents which modulate neovascularization when administered to a subject. Furthermore, known angiogenic agents suffer from a number of limitations. For example, a number of proteins, including thrombospondin 1 (hereinafter "TSP-1"), h-endostatin (hereinafter "endostatin") and
20 h-angiostatin (hereinafter "angiostatin") are thought to have angiogenic activity. However, the cost of producing protein drugs can be prohibitively high. This concern is of a special relevance when a high dose is required in order to achieve therapeutic efficacy.

The full potential of angiogenic therapies is unlikely to be realized until the
25 problems discussed hereinabove have been overcome.

-2-

SUMMARY OF THE INVENTION

This invention is based upon the discovery that angiogenic proteins have a region of about twenty-five amino acids which is substantially conserved among different angiogenic proteins. This region is included in the "angiogenic homology region", also abbreviated as "AHR". It has also been discovered that the angiogenic activity of this class of proteins is attributable, at least in part, to the AHR.

Applicant has also discovered that multivalent ligands comprising peptide derivatives of the AHR of angiogenic proteins modulate angiogenesis in mammals. A multivalent ligand has two or more peptides connected by a covalent bond or linker group to a multilinker backbone. For example, the multivalent ligands Tip-13.40 and Tip-12.40 suppress tumor growth in mice (Example 2). The peptides in Tip-13.40 are represented by SEQ ID NO. 8 and are derivatives of the AHR of TSP-1 (SEQ ID NO.: 1); the peptides in Tip-12.40 are represented by SEQ ID NO.: 7 and are derivatives of the AHR of endostatin (SEQ ID NO.: 2).


A further discovery, reported herein, is that multivalent ligands comprising peptides which are hybrids of the AHR of two different angiogenic proteins also modulate angiogenesis in mammals. For example, the multivalent ligands Tip-14.40, Tip-15.40, Tip-16.40, Tip-18.40 and Tip-19.40 suppress tumor growth in mice (Example 2). The hybrid peptides in Tip-14.40, Tip-15.40, Tip-16.40, Tip-18.40 and Tip-19.40 are represented by SEQ ID NOS.: 3, 4, 10, 11 and 12, respectively, and consist of a subsequence from the AHR of TSP-1 (SEQ ID NO.: 1) and a subsequence from the AHR of endostatin (SEQ ID NO.: 2). In contrast, the multivalent ligand Tip-17.40, which also contains hybrid peptides, stimulates tumor growth in mice. The hybrid peptides in Tip-17.40 are represented by SEQ ID NO.: 9.

Based on the aforementioned discoveries, novel peptides which are peptide derivatives of the AHR of angiogenic peptides are disclosed. Also disclosed are novel peptides which are hybrids of the AHR of angiogenic proteins and peptide

-3-

derivatives of said hybrids. Novel multivalent ligands which comprise these peptide derivatives and/or hybrid peptides and methods of modulating (inhibiting or stimulating) angiogenesis in a subject with these multivalent ligands are other aspects of the present invention.

- 5 One embodiment of the present invention is a multivalent ligand which has angiogenic activity and is represented by Structural Formula (I):



(I)

B is a multilinker backbone.

- 10 n is an integer from two to about twenty.


Each L is a covalent bond or linking group.

Each P is a peptide having from about 10 to about 30 amino acid residues.

- At least two of the peptides are peptide derivatives of an AHR of an angiogenic protein, a hybrid peptide or a peptide derivative of a hybrid peptide. Each P and
15 each linker or covalent bond are independently chosen.

- Another embodiment of the present invention is a polypeptide multivalent ligand having angiogenic activity. A "polypeptide multivalent ligand" is a tandem repeat polypeptide chain in which two or more peptides P are each separated by a peptide spacer. A polypeptide multivalent ligand is represented by Structural
20 Formula (II):

-4-



(II)

m is an integer from zero to about twenty.

Each P is a peptide having from about ten to about forty amino acid residues.

- 5 At least two of the peptides are peptide derivatives of an AHR of an angiogenic protein, a hybrid peptide or a peptide derivative of a hybrid peptide.

Each S is a peptide spacer having from about five to about thirty amino acids.

- 10 Each peptide P and each peptide spacer are independently chosen. The N-terminus and/or C-terminus of the polypeptide multivalent ligand is optionally substituted, as described below.

- Another embodiment of the present invention is a method of modulating (stimulating or inhibiting) angiogenesis in a subject. The method comprises administering a therapeutically effective amount of a multivalent ligand represented by Structural Formula (I) or a polypeptide represented by Structural Formula (II).
- 15

- The multivalent ligands of the present invention can be used to modulate angiogenesis in a subject. Thus, multivalent ligands which inhibit angiogenesis can be used to treat subjects with cancer and other diseases which respond favorably to drugs which inhibit angiogenesis. Multivalent ligands which stimulate angiogenesis can be used to promote wound healing or to stimulate the growth of new blood vessels to bypass, for example, blood vessel occlusions. Multivalent ligands are likely to be less expensive to produce than angiogenic protein drugs. Moreover, multivalent ligands might enable the administration of lower doses in order to achieve therapeutic efficacy, as compared with a univalent peptide chain. In
- 20

-5-

addition, they can have long *in vivo* lifetimes and good biodistribution when administered orally or parenterally.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a list of sequences illustrating the amino acid sequence for AHR
5 of TSP-1 (SEQ ID NO.: 1), endostatin (SEQ ID NO.: 2), TSP-4 (SEQ ID NO.: 5)
and angiostatin (SEQ ID NO.: 6), indicating with underlining the positions where
homology is substantially preserved and indicating with numbering the relative
position of each amino acid in each amino sequence. Included in Figure 1 is the core
region, in which homology is highly conserved among different angiogenic proteins
10 (positions 1-25), and flanking regions (positions -9 to -1 and 26).

Figure 2 is a list of sequences illustrating the amino acid sequence of hybrid
peptides represented by SEQ ID NO.: 3 and by SEQ ID NO.: 4, which are found in
the multivalent ligands Tip-14.40 and Tip-15.40, respectively.

Figure 3 is a list of sequences illustrating the amino acid sequences for
15 peptide derivatives represented by SEQ ID NO.: 7, SEQ ID NO.: 8, SEQ ID NO.: 9,
SEQ ID NO.: 10, SEQ ID NO.: 11 and SEQ ID NO.: 12, which are found in the
multivalent ligands Tip-12.40, Tip-13.40, Tip-17.40, Tip- 16.40, Tip-18.40 and Tip-
19.40, respectively.

Figure 4A is an illustration showing the structure of a multivalent ligand with
20 an asymmetrical cascading polylysine backbone; Figure 4B is an illustration
showing a multivalent ligand with a symmetrical cascading polylysine backbone;
Figure 4C is an illustration showing a multivalent ligand with a pennant polypeptide
backbone; and Figure 4D is an illustration showing a multivalent ligand with a
pennant polylysine backbone.

25 Figure 5 is the consensus sequence for the angiogenic homology region of
angiogenic proteins.

-6-

Figure 6 is a graph showing the volumes of B16 melanoma tumors in mice (measured in mm³) over time (measured in days) after initiation of treatment with the multivalent ligand Tip-12.40. The tumor volumes in Tip-12.40 treated mice are compared with tumor volumes in control mice (vehicle only).

5 Figure 7 is a graph showing the volumes of B16 melanoma tumors in mice (measured in mm³) over time (measured in days) after initiation of treatment with the multivalent ligand Tip-13.40. The tumor volumes in Tip-13.40 treated mice are compared with tumor volumes in control mice (vehicle only).

10 Figure 8 is a graph showing the volumes of B16 melanoma tumors in mice (measured in mm³) over time (measured in days) after initiation of treatment with the multivalent ligand Tip-16.40 or 17.40. The tumor volumes in Tip-16.40 and Tip-17.40 treated mice are compared with tumor volumes in control mice (vehicle only).

15 Figure 9 is a graph showing the volumes of B16 melanoma tumors in mice (measured in mm³) over time (measured in days) after initiation of treatment with the multivalent ligand Tip-18.40 or Tip-19.40. The tumor volumes in Tip-18.40 and Tip-19.40 treated mice are compared with tumor volumes in control mice (vehicle only).

20 Figure 10 is a graph showing the effect of increasing concentrations of Tip-18.40, measured in µg/ml, on bovine aortic endothelial cells proliferation *in vitro*, measured as the percentage treated cells growing in culture relative to untreated cells.

25 Figure 11 is a graph showing the effect of increasing concentrations of Tip-19.40, measured in µg/ml, on bovine aortic endothelial cells proliferation *in vitro*, measured as the percentage treated cells growing in culture relative to untreated cells.

Figure 12 is a diagram showing a polypeptide multivalent ligand represented by SEQ ID NO.: 15.

-7-

DETAILED DESCRIPTION OF THE INVENTION

“Angiogenic agents”, modulate the development of new blood vessels in mammals. Included within the meaning of the term “angiogenic agents” are molecules which inhibit angiogenesis (“anti-angiogenic agents”) and molecules which
5 stimulate or promote angiogenesis (“pro-angiogenic agents”). The development of new blood vessels in mammals is also referred to as “neovascularization”.

Angiogenic agents include proteins and peptides which modulate angiogenesis, referred to herein as “angiogenic proteins” and “angiogenic peptides”, respectively.

Angiogenic agents also include molecules which induce physiological
10 changes in a mammal which are characteristic of angiogenesis modulation. Examples include modulation (promotion or suppression) of tumor growth, tissue repair and tissue remodelling. Peptides which modulate tumor growth when incorporated into multivalent ligands are considered to be angiogenic. Also included within the definition of angiogenic agents are molecules which modulate cellular
15 processes involved in the genesis of blood vessels or the expression of endothelial cell phenotypes. Examples include endothelial cell proliferation, endothelial cell survival, endothelial cell motility, binding to endothelial cells. Agents which modulate cell functions involved in angiogenesis or expression of endothelial cell phenotypes can be identified by *in vitro* assays which assess, for example,
20 endothelial cell proliferation, endothelial cell survival under restrictive conditions, endothelial cell migration and the binding of white blood cells or platelets to endothelial cells. *In vitro* assays for identifying angiogenic agents are described in Example 3 of the present application, Tolsma, *et al.*, *J. Cell Biol.* 122:497 (1993) and Vogel *et al.*, *J. Cell. Biochem.* 53:74 (1993). The entire teachings of Tolsma
25 and Vogel are incorporated herein by reference.

As noted above, Applicant has discovered that angiogenic proteins have an “AHR”, which includes a core region in which the amino acid sequence is

-8-

substantially conserved among several members. Thus, the amino acid sequence of the AHR core of one angiogenic protein is substantially homologous to the AHR core of other angiogenic proteins. An AHR core has between about 10 to about 30 amino acid residues, generally about twenty-five amino acid residues. Once an
5 AHR is identified, its boundaries can optionally be extended to include from about zero to nine flanking amino acids, preferably zero to about four. A consensus sequence for the AHRs of angiogenic proteins is shown in Figure 5. It has also been found that small peptides having an amino acid sequence corresponding to the amino acid sequence of an AHR can modulate angiogenesis. The location of an AHR is
10 not restricted to any particular part of an angiogenic protein. For example, the AHR of endostatin is located closer to the N-terminus, whereas the AHR of TSP-1 is located near the C-terminus.

“Substantial homology” exists between two amino acid sequences when a sufficient number of amino acid residues at corresponding positions of each amino
15 acid sequence are either identical or structurally related such that a peptide having the first amino acid sequence and a peptide having the second amino acid sequence exhibit similar biological activities. A protein has an AHR when the amino acid sequence of said protein has a subsequence which is substantially homologous to the amino acid sequence of an AHR of an angiogenic protein (or to the consensus
20 sequence for the AHR of angiogenic proteins shown in Figure 5) such that a peptide having an amino acid sequence corresponding to said subsequence (or said consensus sequence) modulates angiogenesis. Generally, there is substantial homology among the amino acid sequences of two AHRs when at least 30%, and preferably at least 40% of the amino acids in one AHR are identical to or structurally
25 related to the amino acid residues in the other AHR. Substantial homology exists between the amino acid sequence of a peptide and the amino acid sequence of an AHR (or the consensus sequence shown in Figure 5) when a sufficient number of amino acids at corresponding positions in the amino acid sequence of the peptide

-9-

and AHR (or consensus sequence) are identical or structurally related such that the peptide is angiogenic. Generally, there is substantial homology between a peptide and an AHR when at least 40%, preferably at least 50% of the amino acids in the peptide are identical to or structurally related to the amino acid residues in the
5 corresponding positions in the AHR, or an angiogenic subsequence thereof.
"Structurally related" is defined hereinbelow.

One embodiment of the present invention is a peptide derivative of an AHR of an angiogenic protein. The peptide derivative has angiogenic activity. Examples include peptide derivatives of a peptide represented by SEQ ID NOS.: 2, 5 or 6.

10 A "peptide derivative of an AHR" includes a peptide having the amino acid sequence of the AHR. A "peptide derivative of an AHR" also includes a peptide having a sequence corresponding to an angiogenic fragment of the AHR. An "angiogenic fragment" is defined to be a peptide whose amino acid sequence corresponds to a subsequence of an AHR, referred to as an "angiogenic
15 subsequence". A subsequence is a sequence of contiguous amino acid residues found within a larger sequence. An angiogenic fragment generally has from about ten to about thirty amino acid residues.

A "peptide derivative" also includes a peptide having a "modified sequence" in which one or more amino acid residues in the original sequence or subsequence
20 have been substituted with a naturally occurring amino acid residue or amino acid residue analog (also referred to as a "modified amino acid residue"). Suitable peptide derivatives have modified sequences which are substantially homologous to the amino acid sequence of an AHR or to an angiogenic subsequence of an AHR. Suitable peptide derivatives also include peptides which are substantially
25 homologous to the consensus sequence for the AHRs of angiogenic proteins, shown in Figure 5. Peptide derivatives generally have between about ten and about thirty amino acid residues.

-10-

In one aspect of the present invention, a peptide derivative has an amino acid sequence corresponding to an angiogenic subsequence of an AHR with between about ten and about fifteen amino acid residues. Zero, one, two or three amino acid residues in the peptide derivative can differ from the amino acid residue(s) in the corresponding position of the subsequence of the AHR. For example, if the subsequence is [AA₁]-[AA₂]-AA₃]-[AA₄]-[AA₅]-[AA₆]-[AA₇]-[AA₈]-[AA₉]-[AA₁₀] and one amino acid residue in the sequence of the peptide derivative differs from the amino acid residue in the corresponding position of the subsequence, then the peptide derivative can be [AA₁']-[AA₂]-AA₃]-[AA₄]-[AA₅]-[AA₆]-[AA₇]-[AA₈]-[AA₉]-[AA₁₀], [AA₁]-[AA₂']-AA₃]-[AA₄]-[AA₅]-[AA₆]-[AA₇]-[AA₈]-[AA₉]-[AA₁₀], [AA₁]-[AA₂]-AA₃']-[AA₄]-[AA₅]-[AA₆]-[AA₇]-[AA₈]-[AA₉]-[AA₁₀], [AA₁]-[AA₂]-AA₃]-[AA₄']-[AA₅]-[AA₆]-[AA₇]-[AA₈]-[AA₉]-[AA₁₀], [AA₁]-[AA₂]-AA₃]-[AA₄]-[AA₅']-[AA₆]-[AA₇]-[AA₈]-[AA₉]-[AA₁₀], [AA₁]-[AA₂]-AA₃]-[AA₄]-[AA₅]-[AA₆']-[AA₇]-[AA₈]-[AA₉]-[AA₁₀], [AA₁]-[AA₂]-AA₃]-[AA₄]-[AA₅]-[AA₆]-[AA₇']-[AA₈]-[AA₉]-[AA₁₀], [AA₁]-[AA₂]-AA₃]-[AA₄]-[AA₅]-[AA₆]-[AA₇]-[AA₈']-[AA₉]-[AA₁₀], [AA₁]-[AA₂]-AA₃]-[AA₄]-[AA₅]-[AA₆]-[AA₇]-[AA₈]-[AA₉']-[AA₁₀] and [AA₁]-[AA₂]-AA₃]-[AA₄]-[AA₅]-[AA₆]-[AA₇]-[AA₈]-[AA₉]-[AA₁₀'], wherein [AA'] is a naturally occurring or modified amino acid different from [AA].

In another aspect of the present invention, a peptide derivative has an amino acid sequence corresponding to an amino acid sequence or to an angiogenic subsequence of an AHR with between about sixteen and about twenty-eight amino acid residues. Zero, one, two, three or four amino acid residues in the peptide derivative can differ from the amino acid residue(s) in the corresponding position of the sequence or subsequence of the AHR.

An "amino acid residue" is a moiety found within a peptide and is represented by -NH-CHR-CO-, wherein R is the side chain of a naturally occurring amino acid. When referring to a moiety found within a peptide, the terms "amino acid residue" and "amino acid" are used interchangeably in this application. An

-11-

“amino acid residue analog” includes D or L configurations having the following formula: -NH-CHR-CO-, wherein R is an aliphatic group, a substituted aliphatic group, a benzyl group, a substituted benzyl group, an aromatic group or a substituted aromatic group and wherein R does not correspond to the side chain of a naturally-
5 occurring amino acid.

Suitable substitutions for amino acid residues in the sequence of an AHR or an angiogenic subsequence of an AHR include conservative substitutions which result in peptide derivatives which are angiogenic agents. A conservative substitution is a substitution in which the substituting amino acid (naturally
10 occurring or modified) is structurally related to the amino acid being substituted, i.e., has about the same size and electronic properties as the amino acid being substituted. Thus, the substituting amino acid would have the same or a similar functional group in the side chain as the original amino acid.

A “conservative substitution” also refers to utilizing a substituting amino acid
15 which is identical to the amino acid being substituted except that a functional group in the side chain is functionalized with a suitable protecting group. Suitable protecting groups are described in Green and Wuts, *“Protecting Groups in Organic Synthesis”*, John Wiley and Sons, Chapters 5 and 7, 1991, the teachings of which are incorporated herein by reference. Preferred protecting groups are those which
20 facilitate transport of the peptide through membranes, for example, by reducing the hydrophilicity and increasing the lipophilicity of the peptide, and which can be cleaved, either by hydrolysis or enzymatically (Ditter *et al.*, *J. Pharm. Sci.* 57:783 (1968); Ditter *et al.*, *J. Pharm. Sci.* 57:828 (1968); Ditter *et al.*, *J. Pharm. Sci.* 58:557 (1969); King *et al.*, *Biochemistry* 26:2294 (1987); Lindberg *et al.*, *Drug Metabolism and Disposition* 17:311 (1989); Tunek *et al.*, *Biochem. Pharm.* 37:3867 (1988), Anderson *et al.*, *Arch. Biochem. Biophys.* 239:538 (1985) and Singhal *et al.*, *FASEB J.* 1:220 (1987)). Suitable hydroxyl protecting groups include ester, carbonate and carbamate protecting groups. Suitable amine protecting groups
25

-12-

include acyl groups and alkoxy or aryloxy carbonyl groups, as described above for N-terminal protecting groups. Suitable carboxylic acid protecting groups include aliphatic, benzyl and aryl esters, as described below for C-terminal protecting groups. In one embodiment, the carboxylic acid group in the side chain of one or
5 more glutamic acid or aspartic acid residues in a peptide of the present invention is protected, preferably as a methyl, ethyl, benzyl or substituted benzyl ester, more preferably as a benzyl ester.

Provided below are groups of naturally occurring and modified amino acids in which each amino acid in a group has similar electronic and steric properties.
10 Thus, a conservative substitution can be made by substituting an amino acid with another amino acid from the same group. It is to be understood that these groups are non-limiting, i.e. that there are additional modified amino acids which could be included in each group.

Group I includes leucine, isoleucine, valine, methionine and modified amino
15 acids having the following side chains: ethyl, *n*-propyl *n*-butyl. Preferably, Group I includes leucine, isoleucine, valine and methionine.

Group II includes glycine, alanine, valine and a modified amino acid having an ethyl side chain. Preferably, Group II includes glycine and alanine.

Group III includes phenylalanine, phenylglycine, tyrosine, tryptophan,
20 cyclohexylmethyl, and modified amino residues having substituted benzyl or phenyl side chains. Preferred substituents include one or more of the following: halogen, methyl, ethyl, nitro, -NH₂, methoxy, ethoxy and -CN. Preferably, Group III includes phenylalanine, tyrosine and tryptophan.

-13-

Group IV includes glutamic acid, aspartic acid, a substituted or unsubstituted aliphatic, aromatic or benzylic ester of glutamic or aspartic acid (e.g., methyl, ethyl, *n*-propyl *iso*-propyl, cyclohexyl, benzyl or substituted benzyl), glutamine, asparagine, -CO-NH-alkylated glutamine or asparagine (e.g., methyl, ethyl, *n*-propyl and *iso*-propyl) and modified amino acids having the side chain $-(CH_2)_3-COOH$, an ester thereof (substituted or unsubstituted aliphatic, aromatic or benzylic ester), an amide thereof and a substituted or unsubstituted N-alkylated amide thereof. Preferably, Group IV includes glutamic acid, aspartic acid, methyl aspartate, ethyl aspartate, benzyl aspartate and methyl glutamate, ethyl glutamate and benzyl glutamate, glutamine and asparagine.

Group V includes histidine, lysine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline and 2-amino-4-guanidinobutanoic acid, homologs of lysine, homologs of arginine and homologs of ornithine. Preferably, Group V includes histidine, lysine, arginine and ornithine. A homolog of an amino acid includes from 1 to about 3 additional or subtracted methylene units in the side chain.

Group VI includes serine, threonine, cysteine and modified amino acids having C1-C5 straight or branched alkyl side chains substituted with -OH or -SH, for example, $-CH_2CH_2OH$, $-CH_2CH_2CH_2OH$ or $-CH_2CH_2OHCH_3$. Preferably, Group VI includes serine, cysteine or threonine.

In another aspect, suitable substitutions for amino acid residues in the sequence of an AHR or an angiogenic subsequence of an AHR include "severe" substitutions which result in peptide derivatives which are angiogenic agents.

Severe substitutions which result in peptide derivatives that are angiogenic agents

-14-

are much more likely to be possible in positions which are not highly conserved in the AHRs of angiogenic proteins than at positions which are highly conserved.

Figure 1 shows the amino acid sequence of TSP-1, TSP-4, endostatin and angiostatin and indicates with underlines the positions which are highly conserved.

5 A "severe substitution" is a substitution in which the substituting amino acid (naturally occurring or modified) has significantly different size and/or electronic properties compared with the amino acid being substituted. Thus, the side chain of the substituting amino acid can be significantly larger (or smaller) than the side chain of the amino acid being substituted and/or can have functional groups with
10 significantly different electronic properties than the amino acid being substituted. Examples of severe substitutions of this type include the substitution of phenylalanine or cyclohexylmethyl glycine for alanine, isoleucine for glycine, a D amino acid for the corresponding L amino acid or $-\text{NH}-\text{CH}[(\text{-CH}_2)_5\text{-COOH}]-\text{CO}-$ for aspartic acid. Alternatively, a functional group may be added to the side chain,
15 deleted from the side chain or exchanged with another functional group. Examples of severe substitutions of this type include adding an amine or hydroxyl, carboxylic acid to the aliphatic side chain of valine, leucine or isoleucine, exchanging the carboxylic acid in the side chain of aspartic acid or glutamic acid with an amine or deleting the amine group in the side chain of lysine or ornithine. In yet another
20 alternative, the side chain of the substituting amino acid can have significantly different steric and electronic properties that the functional group of the amino acid being substituted. Examples of such modifications include tryptophan for glycine, lysine for aspartic acid and $-(\text{CH}_2)_4\text{COOH}$ for the side chain of serine. These examples are not meant to be limiting.

25 A "hybrid peptide", as used herein, is an angiogenic peptide having from about 10 to about 30 amino acid residues. Each amino acid residue (naturally occurring or modified) in the amino acid sequence of a hybrid peptide is: 1) identical to the amino acid residue at the corresponding position in the AHR of a first

-15-

angiogenic protein; 2) an amino acid residue structurally related thereto; 3) identical to the amino acid residue at the corresponding position in the AHR of a second angiogenic protein; or 4) an amino acid residue structurally related thereto.

As noted above, replacing an amino acid residue with a structurally related amino acid residue is referred to as a "conservative substitution".

One embodiment of the present invention is a "TSP-1/endostatin hybrid peptide". Each amino acid residue of the TSP-1/endostatin hybrid is identical to or a conservative substitution of the amino acid residue at the corresponding position of TSP-1 or endostatin. In one specific example, the TSP-1/endostatin hybrid peptide is a 25-mer in which the amino acid residue at each position of the amino acid sequence is identical to the amino acid residue at the corresponding position of TSP-1 or endostatin. In another specific example, the serine residue at position seven of the 25-mer hybrid peptide described above is replaced with a threonine residue. As can be seen from Figure 1, a serine residue is present at position seven of the amino acid sequence of the AHR of both TSP-1 and endostatin. Thus, replacement of serine with threonine represents a conservative substitution. Other specific examples include fragments of the 25-mer peptides described above having at least ten amino acid residues.

Another embodiment of the present invention is a hybrid peptide having a sequence of amino acids AA₁ through AA₂₈ or a subsequence thereof having at least ten amino acids, wherein:

AA₁ is lysine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, glycine or alanine;

AA₂ is aspartic acid, asparagine, glutamic acid, glutamine, leucine, valine, isoleucine, methionine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

-16-

AA₃ is phenylalanine, alanine, tyrosine, tryptophan, leucine, isoleucine, methionine, valine or glycine;

AA₄ is threonine, glycine, alanine, cysteine or serine;

AA₅ is alanine, threonine, glycine, cysteine or serine;

5 AA₆ is phenylalanine, tyrosine or tryptophan;

AA₇ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid;

AA₈ tryptophan, alanine, phenylalanine, tyrosine or glycine;

10 AA₉ is arginine, phenylalanine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, lysine, ornithine, tyrosine or tryptophan;

AA₁₀ is leucine, isoleucine, methionine or valine;

AA₁₁ is serine, threonine or alanine;

AA₁₂ is histidine, serine, threonine, cysteine, lysine or ornithine;

15 AA₁₃ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₄ is proline, leucine, valine, isoleucine or methionine;

20 AA₁₅ is lysine, glutamine, histidine, ornithine, asparagine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₆ is threonine, aspartic acid, serine, glutamic acid or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

AA₁₇ is glycine, leucine, alanine, valine, isoleucine or methionine;

AA₁₈ is phenylalanine, tyrosine or tryptophan;

25 AA₁₉ is isoleucine, serine, valine, leucine, methionine, cysteine or threonine;

-17-

AA₂₀ is arginine, isoleucine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline, 2-amino-4-guanidinobutanoic acid, leucine, valine or methionine;

AA₂₁ is methionine, isoleucine, leucine or valine;

5 AA₂₂ valine, arginine, leucine, isoleucine, methionine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

10 AA₂₃ methionine, arginine, leucine, isoleucine, valine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₂₄ is phenylalanine, alanine, tyrosine, tryptophan or glycine;

AA₂₅ is aspartic acid, asparagine, glutamic acid, glutamine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

15 AA₂₆ is glycine, arginine, alanine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

20 AA₂₇ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid; and

AA₂₈ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid.

25 The hybrid peptide has angiogenic activity and is not represented by SEQ ID NO.: 1 or a subsequence thereof.

Other embodiments include hybrid peptides in which each amino acid residue of the hybrid peptide is identical to or a conservative substitution of the amino acid residue at the corresponding position of the AHR of: TSP-1 or TSP-4;

-18-

TSP-1 or angiostatin; endostatin or TSP-4; endostatin or angiostatin; TSP-4 or angiostatin and TSP-1 or endostatin.

Preferably, the amino acid sequence of a hybrid peptide comprises a subsequence from a first AHR and a subsequence from a second AHR. Each
5 subsequence has from about five to about fifteen amino acid residues. In one specific example, the first seven amino acid residues in TSP-1/endostatin hybrid correspond to a subsequence consisting of the first seven amino acid residues of the AHR of TSP-1; the last ten amino acid residues in the TSP-1/endostatin hybrid correspond to a subsequence consisting of the last ten amino acid residues of the
10 AHR of endostatin; and amino acids eight through fourteen are identical to or a conservative substitution of the amino acid residue at the corresponding position of TSP-1 or endostatin.

More preferably, the amino acid sequence of a hybrid peptide consists of a subsequence from a first AHR and a subsequence from a second AHR. Each
15 subsequence has from about five to about fifteen amino acid residues. The two subsequences can be equal in length, e.g., both subsequences can be 9-mers or 12-mers. Alternatively, the two subsequences can be of different lengths, e.g., a 12-mer and a 13-mer. One example, a peptide represented by SEQ ID NO.: 3, is a 22-mer in which the first eleven amino acids correspond to a subsequence of the first eleven
20 amino acid residues of the AHR of endostatin and the second eleven amino acids correspond to a subsequence of the second eleven amino acid residues of the AHR of TSP-1. Another example, a peptide represented by SEQ ID NO.: 4, is a 22-mer in which the first eleven amino acids correspond to a subsequence of the first eleven amino acid residues of the AHR of TSP-1 and the second eleven amino acids
25 correspond to a subsequence of the second eleven amino acid residues of the AHR of endostatin.

Another embodiment of the present invention is a peptide derivative of a hybrid peptide, for example, a peptide derivative of a peptide represented by SEQ ID

-19-

NO.: 3, SEQ ID NO.: 4 or SEQ ID NO.: 9-12. Included within the definition of "peptide derivative of a hybrid peptide" are angiogenic fragments of hybrid peptides, which generally have at least about ten amino acid residues. A "peptide derivative of a hybrid peptide" also includes a peptide having a "modified sequence" in which one or more amino acids in the hybrid peptide have been substituted with a naturally occurring amino acid or amino acid analog (also referred to as a "modified amino acid"). Suitable modified sequences are those which are substantially homologous to the amino acid sequence of the hybrid peptide or to an angiogenic subsequence there. Peptide derivatives generally have between about ten and about thirty amino acid residues.

In one aspect of the present invention, a peptide derivative of a hybrid peptide, e.g., SEQ ID NOS.: 3, 4 or 9-12 has an amino acid sequence corresponding to an angiogenic subsequence of the hybrid peptide with between about ten and about fifteen amino acid residues. Zero, one, two or three amino acid residues in the peptide derivative can differ from the amino acid residue(s) in the corresponding position of the subsequence of the hybrid peptide.

In another aspect of the present invention, a peptide derivative has an amino acid sequence corresponding to the amino acid sequence of the hybrid peptide or to an angiogenic subsequence thereof with between about sixteen and about twenty-eight amino acid residues. Zero, one, two, three or four amino acid residues in the peptide derivative can differ from the amino acid residue(s) in the corresponding position of the sequence or subsequence of the hybrid peptide.

Optionally, the C-terminus or the N-terminus of the peptides of the present invention, or both, can be substituted with a carboxylic acid protecting group or an amine protecting group, respectively. Suitable protecting groups are described in Green and Wuts, *"Protecting Groups in Organic Synthesis"*, John Wiley and Sons, Chapters 5 and 7, 1991, the teachings of which are incorporated herein by reference. As with protecting groups for functional groups found in the amino acid residue side

-20-

chains, preferred protecting groups are those which facilitate transport of the peptide into a cell, for example, by reducing the hydrophilicity and increasing the lipophilicity of the peptide. Examples of N-terminal protecting groups include acyl groups (-CO-R₁) and alkoxy carbonyl or aryloxy carbonyl groups (-CO-O-R₁),

5 wherein R₁ is an aliphatic, substituted aliphatic, benzyl, substituted benzyl, aromatic or a substituted aromatic group. Specific examples of acyl groups include acetyl, (ethyl)-CO-, *n*-propyl-CO-, *iso*-propyl-CO-, *n*-butyl-CO-, *sec*-butyl-CO-, *t*-butyl-CO-, phenyl-CO-, substituted phenyl-CO-, benzyl-CO-, (substituted benzyl)-CO- and myristoyl. Examples of alkoxy carbonyl and aryloxy carbonyl groups include

10 CH₃-O-CO-, (ethyl)-O-CO-, *n*-propyl-O-CO-, *iso*-propyl-O-CO-, *n*-butyl-O-CO-, *sec*-butyl-O-CO-, *t*-butyl-O-CO-, phenyl-O-CO-, substituted phenyl-O-CO- and benzyl-O-CO-, (substituted benzyl)-O-CO-. The carboxyl group at the C-terminus can be protected, for example, as an amide (i.e., the hydroxyl group at the C-terminus is replaced with -NH₂, -NHR₂ and -NR₂R₃) or ester (i.e. the hydroxyl group

15 at the C-terminus is replaced with -OR₂). R₂ and R₃ are independently an aliphatic, substituted aliphatic, benzyl, substituted benzyl, aryl or a substituted aryl group. In addition, taken together with the nitrogen atom, R₂ and R₃ can form a C4 to C8 heterocyclic ring with from about 0-2 additional heteroatoms such as nitrogen, oxygen or sulfur. Examples of suitable heterocyclic rings include piperidinyl,

20 pyrrolidinyl, morpholino, thiomorpholino or piperazinyl. Examples of C-terminal protecting groups include -NH₂, -NHCH₃, -N(CH₃)₂, -NH(ethyl), -N(ethyl)₂, -N(methyl)(ethyl), -NH(benzyl), -N(C1-C4 alkyl)(benzyl), -NH(phenyl), -N(C1-C4 alkyl)(phenyl), -OCH₃, -O-(ethyl), -O-(*n*-propyl), -O-(*n*-butyl), -O-(*iso*-propyl), -O-(*sec*-butyl), -O-(*t*-butyl), -O-benzyl and -O-phenyl.

25 As used herein, aliphatic groups include straight chained, branched or cyclic C1-C6 hydrocarbons which are completely saturated, which contain one or two heteroatoms such as nitrogen, oxygen or sulfur and/or which contain one or more units of unsaturation. Aromatic groups include carbocyclic aromatic groups such as

-21-

phenyl and naphthyl and heterocyclic aromatic groups such as imidazolyl, indolyl, thienyl, furanyl, pyridyl, pyranyl, pyranol, oxazolyl, benzothienyl, benzofuranyl, quinoliny, isoquinoliny and acridintyl.

Suitable substituents on an aliphatic, aromatic or benzyl group include, for example, -OH, halogen (-Br, -Cl, -I and -F) -O(aliphatic, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group), -CN, -NO₂, -COOH, -NH₂, -NH(aliphatic group, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group), -N(aliphatic group, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group)₂, -COO(aliphatic group, substituted aliphatic, benzyl, substituted benzyl, aryl or substituted aryl group), -CONH₂, -CONH(aliphatic, substituted aliphatic group, benzyl, substituted benzyl, aryl or substituted aryl group)), -SH, -S(aliphatic, substituted aliphatic, benzyl, substituted benzyl, aromatic or substituted aromatic group) and -NH-C(=NH)-NH₂. A substituted benzylic or aromatic group can also have an aliphatic or substituted aliphatic group as a substituent. A substituted aliphatic group can also have a benzyl, substituted benzyl, aryl or substituted aryl group as a substituent. A substituted aliphatic, substituted aromatic or substituted benzyl group can have more than one substituent.

A "multivalent ligand" is a molecule having an array peptides, preferably from two to about twenty peptides. Each peptide is connected to a multilinker backbone, either by a covalent bond or through a linker group. Each peptide derivative and each linker or covalent bond is independently chosen. A multivalent ligand is represented by Structural Formula (I).

A "polypeptide multivalent ligand" is a tandem repeat polypeptide chain in which two or more peptides P are each separated by a peptide spacer. A polypeptide multivalent ligand is represented by Structural Formula (II).

At least two of the peptides P in a multivalent ligand or the polypeptide multivalent ligand of the present invention are hybrid peptides, peptide derivatives

-22-

of hybrid peptides, peptide derivatives of AHRs, or combinations thereof. A multivalent ligand or a polypeptide multivalent ligand can also have one or more other peptides P which do not significantly lower its angiogenic activity. Preferably, however, all of the peptides P in a multivalent ligand or a polypeptide multivalent
5 ligand are hybrid peptides, peptide derivatives of hybrid peptides, and/or peptide derivatives of an AHR. Optionally, the C-terminus or the N-terminus of a polypeptide multivalent ligand can include amino acid sequences which might assist in its separation, such as hemagglutinin-antigen or His-Tag. A multivalent ligand or a polypeptide multivalent ligand can have peptides P which are derivatives of
10 different AHRs and/or peptides P which are derivatives of the same AHR but which have different amino acid sequences. A multivalent ligand or a polypeptide multivalent ligand can have peptides P which are hybrids of a different pair of AHRs and/or peptides which are hybrids of the same two AHRs but which have different amino acid sequences. Similarly, a multivalent ligand or a polypeptide multivalent
15 ligand can have peptides P which are derivatives of different hybrid peptides and/or peptides p which are derivatives of the same hybrid peptide but which have different amino acid sequences. More preferably, however, the peptides of a multivalent ligand or a polypeptide multivalent ligand are all the same.

Examples of multivalent ligands are described in the Table. An example of a
20 polypeptide multivalent ligand is shown in Figure 12 and is represented by SEQ ID NO.: 15. Other examples include wherein Tip-19.40 in SEQ ID NO.: 15 is replaced by Tip-12.40, Tip-13.40, Tip-14.40, Tip-15.40, Tip-16.40, Tip-17.40 or Tip-18.40, and/or wherein the polypeptide multivalent ligand has from one to twenty peptide spacers.

25 A multilinker backbone is a linear or branched molecule having a multiplicity of appropriately spaced reactive groups, each of which can react with a functional group in a peptide or linker. Suitable multilinker backbones are biocompatible and, after attachment of the peptide derivatives, are suitable for

-23-


parenteral or oral administration. Generally, the multilinker backbones have molecular weights less than about 20,000 atomic mass units (amu) and typically comprise between two to about a hundred attachment sites. Not all attachment sites need be occupied.

5 Reactive functional groups in a multilinker backbone serve as attachment sites for the peptides or linkers. Attachment sites are "appropriately spaced" when 1) steric hindrance does not substantially interfere with forming covalent bonds between some of the reactive functional groups and the peptide.

 Suitable reactive groups in a multilinker backbone include amines,
 10 carboxylic acids, alcohols, aldehydes and thiols. An amine group in a multilinker backbone can form a covalent bonds with the C-terminal of a peptide derivative or a carboxylic acid functional group in a linker group. A carboxylic acid group or an aldehyde in a multilinker backbone form a covalent bond with the N-terminus of a peptide derivative or an amine group in a linker group. An alcohol group in a
 15 multilinker backbone can form a covalent bond with the C-terminus of a peptide derivative or a carboxylic acid group in a linker group. A thiol group in a multilinker backbone can form a disulfide bond with a cysteine in a peptide derivative or a thiol group in a linker group. Bonds can also be formed between reactive functional groups in the multilinker backbone and appropriate functional
 20 groups in the amino acid side chains of the attached peptides, as described above. The functionality which connects each peptide to the multilinker backbone can be different, but is preferably the same for all peptides.

 Examples of suitable multilinker backbones include polymers with pendant reactive groups such as $[-(\text{CH}_2)_n\text{-CHR}(\text{NH}_2)\text{-}]_m$, $[-\text{CHRNH}_2\text{-}]_m$, $[-(\text{CH}_2)_n\text{-CHR}(\text{COOH})\text{-}]_m$ (R is, for example, an aliphatic or aryl group), polyethylene glycols and polypeptides polyvinyl pyrrolidones and derivatives thereof, and aromatic polymers are represented by, for example, by the following Structural Formula:

-24-



Also included are polyurethane multilinker backbones, examples of which are disclosed in Lin *et al.*, *J. Biomed. Material. Res.* 28:329 (1994), the entire teachings of which are incorporated herein by reference. Polypeptides are preferred multilinker backbones.

Examples of suitable polypeptide multivalent backbones are disclosed in Tam, *Journal of Immunological Methods* 196:17 (1996), the entire teachings of which are incorporated herein by reference. Suitable polypeptide multilinker backbones generally have between about three and about forty amino acid residues. As with other multilinker backbones, they typically have between about two and about twenty attachment sites, which are often functional groups located in the amino acid residue side chains. However, alpha amino groups and alpha carboxylic acids can also serve as attachment sites.

Preferred polypeptide multilinker backbones include polylysines, polyornithines, polycysteines, polyglutamic acid and polyaspartic acid. Optionally, amino acid residues with inert side chains, e.g., glycine, alanine and valine, can be included in the amino acid acid sequence. The polypeptides can be pennant or cascading. A "pennant polypeptide" is linear. As with polypeptides typically found in nature, the amide bonds of a pennant polypeptide are formed between the alpha

-25-

amine of one amino acid residue and the alpha carboxylic acid of the next amino acid residue. When n is less than five, there are typically 0-6 amino acids between attachment sites; when n is greater than five, there are typically 1-6 amino acids between attachment sites. A "cascading polypeptide" is branched with at least some of the amide bonds formed between the side chain functional group of one amino acid residue and the alpha amino group or alpha carboxylic acid group of the next amino acid residue. For example, at least some of the amide bonds of a cascading polylysine are formed between the epsilon amine group of a lysine residue and the carboxylic acid residue of the next lysine residue. Examples of cascading and pennant polylysines are shown in Figure 4.

Suitable linkers are inert groups which can connect a peptide derivative to a multilinker backbone. In one example, the linker is an oligopeptide of from about one to about 10 amino acids consisting of amino acids with inert side chains. Suitable oligopeptide include polyglycine, polyserine, polyproline, polyalanine and oligopeptides consisting of alanyl and/or serinyl and/or prolanyl and/or glycyl amino acid residues. In another example, the linker is $X_1-(CH_2)_m-X_2$ or X_1 -polyethylene-glycol- X_2 . X_1 and X_2 are the residues of a functional group which is connected by a covalent bond to a suitable functional group residue in the multilinker backbone or peptide derivative, respectively. Examples of X_1 and X_2 include: 1) the residue of an alcohol group which forms an ester with the residue of a carboxylic acid group in the multilinker backbone or peptide derivative; 2) the residue of an amine group which forms an amide with the residue of a carboxylic acid group in the multilinker backbone or peptide derivative; 4) the residue of a carboxylic acid or aldehyde group which forms an amide with the residue of an amine in the multilinker backbone or peptide derivative; or 5) the residue of a thiol group which forms a disulfide with the residue of a thiol group in the multilinker backbone or peptide derivative. m is an integer from two to about 20.

-26-

The peptides in a multivalent ligand can be connected to the multilinker backbone by covalent bonds, linker groups or a combination thereof. The linking groups can be the same or different. Preferably, every peptide in a multivalent ligand is connected to the multilinker backbone by a covalent bond. Alternatively,
5 every peptide in a multivalent ligand is connected to the multilinker backbone by the same linking group, e.g., a glycine residue or a glycyl-glycyl dipeptide.

Examples of multivalent ligands are referred to herein as Tip-12.40, Tip-13.40, Tip-14.40, Tip-15.40, Tip-16.40, Tip-17.40, Tip-18.40 and Tip-19.40. These multivalent ligands are four-branch penannt polylysine trimers, indicating that four
10 peptides are attached to a linear trimeric polypeptide backbone consisting of three lysine residues. The attachment is by means of a peptide bond between the C-terminus of each peptide or linking oligopeptide and the amino group in one of the three lysine side chains or the N-terminus of the polylysine. All peptides and all linkers in a given multivalent ligand are the same. Further structural information is
15 provided below in the Table:

-27-

Table

Multivalent Ligand	Sequence of Each Peptide	Linking Group
Tip-12.40	Represented by SEQ ID NO.: 7	Covalent Bond
Tip-13.40	Represented by SEQ ID NO.: 8	Covalent Bond
Tip-14.40	Represented by SEQ ID NO.: 3	Covalent Bond
Tip-15.50	Represented by SEQ ID NO.: 4	Covalent Bond
Tip-16.40	Represented by SEQ ID NO.: 10	-Gly-Gly-
Tip-17.40	Represented by SEQ ID NO.: 9	-Gly-Gly-
Tip-18.40	Represented by SEQ ID NO.: 11	-Gly-Gly-
Tip-19.40	Represented by SEQ ID NO.: 12	-Gly-Gly-

A polypeptide spacer S, shown in Structural Formula (II), is a peptide having from about five to about forty amino acid residues. The spacers in a polypeptide multivalent ligand are independently chosen, but are preferably all the same. The spacers should allow for flexibility of movement in space for the flanking peptides P, and are therefore typically rich in small amino acids, for example, glycine, serine, proline or alanine. Preferably, peptide spacers contain at least 60%, more preferably

-28-

at least 80% glycine or alanine. In addition, peptide spacers generally have little or no biological and antigenic activity. Preferred spacers are $(\text{Gly-Pro-Gly-Gly})_x$ and $(\text{Gly}_4\text{-Ser})_y$, wherein x is an integer from about three to about nine and y is an integer from about one to about eight. Specific examples of suitable spacers include $(\text{Gly}_4\text{-Ser})_3$ (SEQ ID NO.: 13) or $(\text{Gly}_4\text{-Ser})_4$ (SEQ ID NO.: 14). Spacers can also include from one to about four amino acids which create a restriction site, but which are non-angiogenic and which are not biologically active.

The multivalent ligands and polypeptide multivalent ligands of the present invention are angiogenic agents and can therefore be used to modulate angiogenesis in a subject or individual. A "subject" is preferably a human, but can also be animals in need of treatment, e.g., veterinary animals (e.g., dogs, cats, and the like), farm animals (e.g., cows, pigs, horses and the like) and laboratory animals (e.g., rats, mice, guinea pigs and the like).

Angiogenic agents, including the multivalent ligands and polypeptide multivalent ligands disclosed herein, can be used to treat a wide variety of disease conditions, including cancer, cardiovascular diseases (e.g., arteriosclerosis, ischaemic limbs and ischaemic myocardium) obesity, osteoarthritis, duodenal ulcers, and abnormal ocular neovascularization associated, for example, with diabetes mellitus.

A "therapeutically effective amount" is the quantity of compound which results in an improved clinical outcome as a result of the treatment compared with a typical clinical outcome in the absence of the treatment. Alternatively, an "improved clinical outcome" includes a longer life expectancy for subjects with the disease as a result of the treatment or results in the subject with the disease experiencing fewer symptoms or complications of the disease as a result of the treatment. With respect to cancer, an "improved clinical outcome" includes a longer life expectancy. Alternatively, it can include slowing or arresting the rate of growth of a tumor, causing a shrinkage in the size of the tumor, a decreased rate of metastasis and/or

-29-

improved quality of life (e.g., a decrease in physical discomfort or an increase in mobility).

With respect to abnormal ocular neovascularization associated, for example, with diabetes, "improved clinical outcome" refers to slowing, retarding or reversing
5 the loss of sight associated with the disease. Alternatively, it refers to a reduction in retinopathy.

With respect to obesity, an improved clinical outcome refers to increased weight reduction per calory intake. Alternatively, it refers to a decrease in the complications which are a consequence of obesity, for example heart disease such as
10 arteriosclerosis and high blood pressure.

With respect to osteoarthritis, an improved clinical outcome refers to slowing, arresting or reversing the degradation and loss of function typically observed in a joint afflicted with osteoarthritis, e.g. by reducing the rate of cartilage degradation in the joint. Alternatively, it refers to lessening the pain and/or
15 inflammation associated with osteoarthritis.

With respect to wound healing, an improved clinical outcome can refer to a more rapid rate of wound closure, less wound contraction and/or less scarring.

With respect to neovascularization to bypass occluded blood vessels, a "therapeutically effective amount" is a quantity which results in the formation of new
20 blood vessels which can transport at least some of the blood which normally would pass through the blocked vessel.

The amount of the multivalent ligands or polypeptide multivalent ligands administered to the subject will depend on the type and severity of the disease and on the characteristics of the subject, such as general health, age, sex, body weight
25 and tolerance to drugs. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. Typically, a therapeutically effective amount of the multivalent ligand can range from about 1 mg per kilogram body weight per day to about 1000 mg per kilogram body weight per day for an adult.

-30-

Preferably, the dosage ranges from about 1 mg per kilogram body weight per day to about 100 mg per kilogram body weight per day.

The multivalent ligands and polypeptide multivalent of the present invention can, for example, be administered orally or parenterally. Parenteral administration
5 can include, for example, systemic administration, such as by intramuscular, intravenous, subcutaneous, or intraperitoneal injection.

The multivalent ligands and polypeptide multivalent ligands can be administered to the subject in conjunction with an acceptable pharmaceutical carrier as part of a pharmaceutical composition for treating the diseases discussed above.
10 Suitable pharmaceutical carriers may contain inert ingredients which do not interact with the peptide or peptide derivative. Standard pharmaceutical formulation techniques may be employed such as those described in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA. Suitable pharmaceutical carriers for parenteral administration include, for example, sterile water, physiological saline,
15 bacteriostatic saline (saline containing about 0.9% mg/ml benzyl alcohol), phosphate-buffered saline, Hank's solution, Ringer's-lactate and the like. Methods for encapsulating compositions (such as in a coating of hard gelatin or cyclodextran) are known in the art (Baker, *et al.*, *Controlled Release of Biological Active Agents*, John Wiley and Sons, 1986).

20 When used to promote wound healing, a multivalent ligand or polypeptide multivalent ligand be applied directly to the wound as part of a pharmaceutical formulation which includes a pharmaceutical carrier. Examples of pharmaceutically acceptable carriers include, for example, commercially available inert gels, or liquids supplemented with albumin, methyl cellulose or a collagen matrix. Typical of such
25 formulations are ointments, creams and gels. Ointments are typically prepared using an oleaginous base, e.g., containing fixed oils or hydrocarbons, such as white petrolatum or mineral oil, or an absorbent base, e.g., consisting of an absorbent anhydrous substance or substances, for example anhydrous lanolin. Following

-31-

formation of the base, the active ingredients are added in the desired concentration. Creams generally comprise an oil phase (internal phase) containing typically fixed oils, hydrocarbons, and the like, such as waxes, petrolatum, mineral oil, and the like, and an aqueous phase (continuous phase), comprising water and any water-soluble substances, such as added salts. The two phases are stabilized by use of an emulsifying agent, for example, a surface active agent, such as sodium lauryl sulfate; hydrophilic colloids, such as acacia colloidal clays, beegum, and the like. Upon formation of the emulsion, the active ingredients are added in the desired concentration. Gels are comprised of a base selected from an oleaginous base, water, or an emulsion-suspension base, as previously described. To the base is added a gelling agent which forms a matrix in the base, increasing its viscosity to a semisolid consistency. Examples of gelling agents are hydroxypropyl cellulose, acrylic acid polymers, and the like. The active ingredients are added to the formulation at the desired concentration at a point preceding addition of the gelling agent.

The multivalent ligands or polypeptide multivalent ligands of the present invention can be co-administered with other pharmaceutically active agents. In one example, the multivalent ligands or polypeptide multivalent ligands are co-administered with other anti-cancer agents when used to treat a subject with cancer.

The peptide derivatives, the multivalent ligands and or polypeptide multivalent ligands of the present invention have many utilities other than for therapy. Some of these uses are discussed in the following paragraphs.

The disclosed multivalent ligands can be used to raise antibodies, both polyclonal and monoclonal, against the peptide derivatives attached thereto.

Methods of raising antibodies against peptide antigens attached to a multilinker backbone are described in Tam, *Proc. Natl. Acad. Sci. USA* 85:5409 (1988), Tam and Lu, *Proc. Natl. Acad. Sci. USA* 86:9084 (1989) and Tam, *Journal of*

-32-

Immunological Methods 196:17 (1996), the entire teachings of which are incorporated herein by reference.

Suitable antibodies can also be raised against the peptide derivatives and hybrid peptides of the present invention by conjugating to a suitable carrier, such as keyhole limpet hemocyanin or serum albumin; polyclonal and monoclonal antibody production can be performed using any suitable technique. A variety of methods for producing monoclonal antibodies have been described (see e.g., Kohler *et al.*, *Nature*, 256: 495-497 (1975) and *Eur. J. Immunol.* 6: 511-519 (1976); Milstein *et al.*, *Nature* 266: 550-552 (1977); Koprowski *et al.*, U.S. Patent No. 4,172,124; Harlow, E. and D. Lane, 1988, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory: Cold Spring Harbor, NY); *Current Protocols In Molecular Biology*, Vol. 2 (Supplement 27, Summer 1994), Ausubel, F.M. *et al.*, Eds., (John Wiley & Sons: New York, NY), Chapter 11, (1991)). Generally, a hybridoma can be produced by fusing a suitable immortal cell line (e.g., a myeloma cell line such as SP2/0) with antibody producing cells. The antibody producing cell, preferably those of the spleen or lymph nodes, can be obtained from animals immunized with the antigen of interest. The fused cells (hybridomas) can be isolated using selective culture conditions, and cloned by limiting dilution. Cells which produce antibodies with the desired specificity can be selected by a suitable assay (e.g., ELISA).

Antibodies, including monoclonal antibodies, against the AHR of angiogenic proteins have a variety of uses. For example, those against or reactive with an angiogenic protein, and preferably which bind specifically to the AHR of said protein, can be used to determine if the angiogenic protein is present in a liquid sample obtained from a subject. The sample is treated with anti-AHR antibody specific for the angiogenic protein. The sample is then analyzed, for example, by Western blotting or immunoprecipitation for complexes between the angiogenic protein and antibody. The sample can be, for example, a cleared lysate of a cell, which is generated for example, by treating cells with a detergent such as sodium

-33-

deoxycholate (0.5% - 1%) or sodium dodecyl sulfate (1%), centrifugating and separating the supernatant from the pellet.

AHRs play a key role in the biological activity of angiogenic proteins, as is evidenced by the fact that multivalent ligands comprising the peptide derivatives or
5 hybrid peptides of the present invention have such a dramatic effect on biological processes such as tumor growth. The polypeptide multivalent ligands, peptide derivatives and hybrid peptides of the present invention can also be used to identify molecules which interact with the AHR of specific angiogenic proteins and whose activities are modulated by them. For example, an affinity column can be prepared
10 to which a specific polypeptide multivalent ligand, peptide derivative or hybrid peptide is covalently attached, directly or via a linker. This column, in turn, can be utilized for the isolation and identification of specific molecules which bind the AHRs of angiogenic proteins and which will also likely bind the angiogenic protein from which the peptide derivative was derived. The molecule can then be eluted
15 from the column, characterized and tested for its ability to interact with angiogenic proteins.

Peptide sequences in the compounds of the present invention may be synthesized by solid phase peptide synthesis (e.g., BOC or FMOC) method, by solution phase synthesis, or by other suitable techniques including combinations of
20 the foregoing methods. The BOC and FMOC methods, which are established and widely used, are described in Merrifield, *J. Am. Chem. Soc.* 88:2149 (1963); Meienhofer, *Hormonal Proteins and Peptides*, C.H. Li, Ed., Academic Press, 1983, pp. 48-267; and Barany and Merrifield, in *The Peptides*, E. Gross and J. Meienhofer, Eds., Academic Press, New York, 1980, pp. 3-285. Methods of solid phase peptide
25 synthesis are described in Merrifield, R.B., *Science*, 232: 341 (1986); Carpino, L.A. and Han, G.Y., *J. Org. Chem.*, 37: 3404 (1972); and Gauspohl, H. *et al.*, *Synthesis*, 5: 315 (1992)). The teachings of these references are incorporated herein by reference.

-34-

Multivalent ligands can be prepared by methods known in the art, including methods disclosed in Tam, *J. of Immunological Methods* 196:17 (1996), Kim *et al.*, *Cancer Research* 54:5005 (1994), Nomizu *et al.*, *Cancer Research* 53:3459 (1993) and Tam, *Proc. Natl. Acad. Sci., USA* 85:5409 (1989), the teachings of which are
5 incorporated herein by reference.

A convenient method of preparing multivalent ligands in a single operation is by stepwise solid phase synthesis, starting with the C-terminus core matrix using a diprotected Boc-Lys(Boc) in Boc chemistry or Fmoc-Lys(Fmoc) in Fmoc chemistry to reach the desired branching. The selected peptide derivative, hybrid peptide or
10 hybrid peptide derivative is then sequentially elongated to the lysinyl core matrix on the resin to form the desired multivalent ligand. This stepwise method produces multivalent ligands with a C → N orientation. Chimeric multivalent ligands having two or more different appended peptides can be also produced in this way by tandemly synthesizing both sequences in a continuous array. Alternatively, different
15 peptides can be synthesized on the different arms of the core matrix, using a core matrix bearing two different amine-protecting groups. Methods to distinguish the α and ϵ -amines of lysines so that different peptides and functional moieties could be introduced have been developed. A common theme in these methods is the manipulation of the orthogonality or differential liability of deprotecting methods.
20 Suitable combinations include: (i) Boc-Fmoc (Tam and Lu, *Proc. Natl. Acad. Sci., USA* 86:9084 (1989); (ii) Fmoc-Dde (Bycroft *et al.*, *J. Chem. Soc. Chem. Commun.* 1993:773 (1993); and (iii) Npys-Fmoc (Ahlborg, *J. Immunol. Methods* 179:269 (1995). The entire teachings of these references are incorporated herein by reference.

25 A polynucleotide encoding a polypeptide multivalent ligand represented by Structural Formula (II) can be prepared by methods disclosed in Lake *et al.*, *Biotechniques* 19:700 (1995), the entire teachings of which are incorporated herein by reference. Alternatively, this gene, encoding the polypeptide multivalent ligand

-35-

represented by Structural Formula (II), can be prepared by synthesizing several separate primers with overlapping regions and then anneal them together, using commonly available automated DNA synthesizers. Following ligation of the annealed polynucleotide, the DNA which code for the desired periodicity of the multivalent ligand can be then selected by its size and its DNA sequence. Once prepared, a polynucleotide encoding a polypeptide multivalent ligand can be expressed using standard recombinant DNA technology. Specifically, the polynucleotide is introduced into a suitable amplifying organism (e.g., *E. coli* or yeast) using an appropriate plasmid vector; the resulting amplifying organism, which contains the polynucleotide, is maintained under conditions suitable for the polynucleotide to be expressed and reproduced by the amplifying organism. Suitable plasmid vectors typically contain an origin of replication (e.g. an autonomously replicating sequence when the plasmid is transformed into yeast), a yeast centromere and a growth origin suitable for growth and replication in the tester cells being transformed. The plasmid vector also contains a gene allowing for selection in the amplifying organism being transformed, e.g. a gene conferring resistance to an antibiotic provided in the growth medium or allowing the expression cells to metabolize an essential nutrient supplied in the growth medium (for example *trp*). Typically, the plasmid also comprises genes suitable for selection and growth in an amplifying organism. Optionally, the plasmid can contain one or more polycloning sites and/or a gene (e.g. *lacZ* gene) which indicates incorporation of the transforming DNA into the plasmid. Plasmids are introduced into the cells of an amplifying organism by methods known in the art, for example, by the calcium phosphate method, electroporation or lithium chloride. Suitable plasmids also have promoter sites which allow the gene being assessed to be efficiently expressed in the amplifying organism (e.g. SV40, *gal 10* or other viral promoters). Specific experimental protocols are disclosed in "Molecular Cloning", by Sambrook J.,

-36-

Fritsch E.F. and Maniatis, T. Cold Spring Harbor Laboratory Press, second edition (1989), the entire teachings of which are incorporated herein by reference.

Other procedures for preparing polypeptide multivalent ligands are disclosed in Röttschlee et al., *Proc. Natl. Acad. Sci. USA* 94:1462 (1997), the entire teachings
5 of which are incorporated herein by reference.

The invention is illustrated by the following examples which are not intended to be limiting in any way.

EXEMPLIFICATION

Example 1 - Preparation of Multivalent Ligands

10 The synthesis of a tetra branched matrix core with an AHR- peptide attached was accomplished manually by a stepwise solid-phase procedure (Merrifield, R.B., *J. Am. Chem. Soc.*, 85:2149-2154 (1963) on *t*-butoxycarbonyl (Boc) β Ala-OCH₂-Pam resin (Mitchell, A.R., *et al.*, *J. Org. Chem.*, 43:2845-2852 (1978) in which 0.05 mmol of β Ala is present in 0.5 g of resin. The synthesis of the first and every
15 subsequent level of the carrier core was achieved using a 4 M excess of preformed symmetrical anhydride of N $^{\alpha}$,N $^{\epsilon}$ -Boc-Lys(Boc) (0.2, 0.4, 0.8 and 1.6 mmol consecutively) in dimethylformamide (HCONMe₂, 12 ml/g resin) followed by a second coupling via dicyclohexylcarbodiimide alone in CH₂Cl₂ to give, after
20 deprotection, the tetra-branched core matrix containing four functional amino groups. The protecting groups for the synthesis of the peptide antigens were Boc groups for the α -amino termini and benzyl alcohol derivatives for most side-chain amino acids. For all residues except arginine, asparagine, glutamine, and glycine, the first coupling for 1 hour, monitored by quantitative ninhydrin test (Sarin, V.K., *et al.*, *Anal. Biochem.*, 117:147-157 (1981) was done with the preformed symmetrical
25 anhydride in CH₂Cl₂, a second coupling in HCONMe₂, and a third (if needed) in *N*-methylpyrrolidone at 50°C (Tam, J. P., in *Proceedings of the Ninth American*

-37-

Peptide Symposium, eds. Deber, C. M., Kopple, K.D. and Hruby, V. J. (Pierce Chem., Rockford, IL) pp. 305-308 (1985)). The coupling of Boc-Asn and Boc-Gln was mediated by the preformed 1-hydroxybenzotriazole ester in HCONMe₂. Boc-Gly and Boc-Arg were coupled with water-soluble dicyclohexylcarbodiimide alone to avoid, respectively, the risk of formation of dipeptide and lactam. To eliminate the polycationic amino groups, which give highly charged macromolecules, the peptide chains were capped on their α -amino group by acetylation in 3 mM acetic anhydride in HCONMe₂ containing 0.3 mmol of *N,N*-dimethylaminopyridine at the completion of the multivalent ligand. The deprotection process was initiated by removing the dinitrophenyl protecting group of His(Dnp) with 1 M thiophenol in HCONMe₂ for 8 hours (3 times and at 50°C if necessary to complete the reaction). The branched peptide oligolysine was removed from the crosslinked polystyrene resin support with the low-high-HF method or the low-high trifluoromethane-sulfonic acid method of cleavage to yield the crude multivalent ligand (85%-93% cleavage yield) (Tam, J.P., *et al.*, *J. Am. Chem. Soc.*, 108:5242-5251 (1986). The crude peptide and resin were then washed with cold ether/mercaptoethanol (99:1, vol/vol., 30 ml) to remove *p*-thiocresol and *p*-cresol, and the peptide was extracted with 100 ml of 8M urea/0.2 M dithiothreitol/0.1 M Tris-HCl buffer, pH 8.0. To remove all the remaining aromatic by-products generated in the cleavage step, the peptide was dialyzed in Spectrum Por 6 tubing, 1000 M, cutoff by equilibration for 24 hours with a deaerated and N₂-purged solution containing 8 M urea, 0.1 M NH₄HCO₃/(NH₄)₂CO₃, pH 8.0, with 0.1 M mercaptoethanol at 0°C for 24 hours. The dialysis was then continued in 8 M and then in 2 M urea-- all in 0.1 M NH₄HCO₃/(NH₄)₂CO₃ buffer, pH 8.0 for 12 hours and then sequentially in H₂O and 1 M HOAc to remove all urea. The multivalent ligand was lyophilized and purified batchwise by high-performance gel-permeation or ion-exchange chromatography. All purified materials were analyzed and found to contain the predicted amino acid sequences.

-38-

Example 2 - Multivalent Ligands Modulate Tumor Growth in Mice

B16 melanoma cells were injected into 7-8 week old female C57BL mice or CBF1 mice, S.C., 2×10^5 cells per mouse, in 0.1 ml volume. One week after the tumor inoculation, the treatment was started by S.C. injection at sites remote from the tumor of the indicated multivalent ligand at a dose 20-30 mg/kg body weight/day for a period of two weeks. Tumors were measured with a dial-caliper and volumes were determined using the formula: $(\text{width})^2 \times \text{length} \times 0.52$. Each group (experimental or control) contained 5-6 mice. Figures 6-9 depict changes in mean tumor-volume over time.

As can be seen from Figures 6-9, Tip-12.40, Tip-13.40, Tip-16.40, Tip-18.40 and Tip-19.40 suppressed tumor growth, whereas Tip-17.40 promoted tumor growth. Tip-14.40 and Tip-15.40 were also tested and showed weak, but statistically significant inhibition of B16 melanoma tumor growth in mice.

Example 3 - Multivalent Ligands Inhibit the Proliferation of Bovine Aortic

Endothelial

Cells In Vitro

The experiments were performed using 96 well, flat bottom, tissue culture microtiter plates. Culture medium was prepared from DMEM, penicillin (100 U/ml), streptomycin (100 $\mu\text{g/ml}$), glutamine (2 mM), 10% endotoxin free bovine calf serum (Hyclone) and 1 ng/ml of basic fibroblast growth factor (any commercial source).

Bovine aortic endothelial cells (referred to herein as "A19 cells") were obtained by procedures disclosed in Gospodorowicz et al., *Proc. Natl. Acad. Sci. USA* 73:4120 (1976), the entire teachings of which are incorporated herein by reference. An A19 cell suspension at 25×10^3 cells/ml was prepared in the above described culture medium and distributed 0.160 ml/well (about 4000 endothelial cells/well).

-39-

Multivalent ligand peptides Tip-18.40 and Tip-19.40 stock solutions was prepared by diluting a 10 mM solution of the HJ peptide in 100% DMSO with phosphate buffered saline (PBS) containing 0.1% BSA. The concentration of HJ peptide in each stock solution was adjusted to nine times the desired concentration
5 of the HJ peptide in the assay mixture.

0.020 ml of each HJ peptide stock solution was added to the corresponding wells about 2 hours after BCE plating, with six replicates for each concentration. In addition, BSA solution with no added HJ peptide was used as a control. The wells were incubated for 72-80 hours at 37°C in a 10% CO₂ humidified incubator.
10 The plates were labeled and the medium discarded. Each plate was then washed one time with PBS (0.200 ml/well). The wells were fixed with 4% formaldehyde PBS (PBS buffered 10% formalin from Fisher Scientific; Catalog No. HC200-1) (0.12 ml/well) for at least 30 minutes.

The wells were washed one time with borate buffer (0.1 M, pH 8.5). Freshly
15 filtered 1% methylene blue solution (0.060 ml/well) was then added to the wells and incubated for 10 minutes at room temperature. The wells were then washed five times with tap water, after which the wells were dried completely. 0.200 ml/well of 0.1 N HCl (0.1 N) was added to extract the color. After extracting overnight, the O.D. was read at 630 nm to determine the relative number of cells per well. The
20 procedure for counting cells is described in greater detail in Oliver et al., *J. of Cell Sci.*, 92:513 (1989), the teachings of which are incorporated herein by reference.

The results are shown in Figures 10 and show that Tip-18.40 and Tip-19.40 both inhibit the proliferation of bovine aortic endothelial cells *in vitro*.

EQUIVALENTS

25 While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without

-40-

departing from the spirit and scope of the invention as defined by the appended claims. Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described specifically herein. Such equivalents are intended to be

5 encompassed in the scope of the claims.

-41-

CLAIMS

What is claimed is:

1. A peptide having the amino acid sequence set forth in SEQ ID NO.: 5, SEQ ID NO.: 6, a subsequence of SEQ ID NO.: 5 having at least ten amino acid residues or a subsequence of SEQ ID NO.: 6 having at least ten amino acid residues.
2. The peptide of Claim 1 wherein the N-terminus is substituted with an amine protecting group and/or the C-terminus is substituted with a carboxylic acid protecting group.
3. The peptide of Claim 1 wherein the peptide has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 5 or SEQ ID NO.: 6 having from ten to fifteen amino acid residues.
4. The peptide of Claim 1 wherein the peptide has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 5 or SEQ ID NO.: 6 having from about ten to about fifteen amino acid residues, wherein one, two or three amino acid residues in the sequence of the peptide can differ from the amino acid residue(s) in the corresponding position(s) of the subsequence.
5. The peptide of Claim 1 wherein the peptide has an amino acid sequence corresponding to SEQ ID NO.: 5, SEQ ID NO.: 6, a subsequence of SEQ ID NO.: 5 having at least sixteen amino acid residues or a subsequence of SEQ ID NO.: 6 having at least sixteen amino acid residues.

-42-

6. The peptide of Claim 1 wherein the peptide has an amino acid sequence corresponding to SEQ ID NO.: 5, SEQ ID NO. 6, a subsequence of SEQ ID NO.: 5 having at least sixteen amino acid residues or a subsequence of SEQ ID NO.: 6 having at least sixteen amino acid residues, wherein one, two, three or four amino acid residues in the sequence of the peptide can differ from the amino acid residue(s) in the corresponding positions of SEQ ID NO.: 5, SEQ ID NO. 6, the subsequence of SEQ ID NO.: 5 or the subsequence of SEQ ID NO.: 6.
7. The peptide of Claim 1 wherein the peptide has an amino acid sequence as set forth in SEQ ID NOS.: 3 or 4.
8. A peptide comprising a sequence of amino acids AA₁ through AA₂₈ or a subsequence thereof having at least ten amino acids, wherein:
- AA₁ is lysine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, glycine or alanine;
- AA₂ is aspartic acid, asparagine, glutamic acid, glutamine, leucine, valine, isoleucine, methionine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;
- AA₃ is phenylalanine, alanine, tyrosine, tryptophan, leucine, isoleucine, methionine, valine or glycine;
- AA₄ is threonine, glycine, alanine, cysteine or serine;
- AA₅ is alanine, threonine, glycine, cysteine or serine;
- AA₆ is phenylalanine, tyrosine or tryptophan;
- AA₇ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid;
- AA₈ tryptophan, alanine, phenylalanine, tyrosine or glycine;

-43-

AA₉ is arginine, phenylalanine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, lysine, ornithine, tyrosine or tryptophan;

AA₁₀ is leucine, isoleucine, methionine or valine;

5 AA₁₁ is serine, threonine or alanine;

AA₁₂ is histidine, serine, threonine, cysteine, lysine or ornithine;

AA₁₃ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₄ is proline, leucine, valine, isoleucine or methionine;

10 AA₁₅ is lysine, glutamine, histidine, ornithine, asparagine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₆ is threonine, aspartic acid, serine, glutamic acid or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

15 AA₁₇ is glycine, leucine, alanine, valine, isoleucine or methionine;

AA₁₈ is phenylalanine, tyrosine or tryptophan;

AA₁₉ is isoleucine, serine, valine, leucine, methionine, cysteine or threonine;

20 AA₂₀ is arginine, isoleucine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline, 2-amino-4-guanidinobutanoic acid, leucine, valine or methionine;

AA₂₁ is methionine, isoleucine, leucine or valine;

25 AA₂₂ valine, arginine, leucine, isoleucine, methionine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₂₃ methionine, arginine, leucine, isoleucine, valine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

-44-

AA₂₄ is phenylalanine, alanine, tyrosine, tryptophan or glycine;

AA₂₅ is aspartic acid, asparagine, glutamic acid, glutamine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

5 AA₂₆ is glycine, arginine, alanine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₂₇ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or
10 2-amino-4-guanidinobutanoic acid; and

AA₂₈ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid.

wherein the peptide has angiogenic activity, with the proviso that the
15 peptide is not represented by SEQ ID NO.: 1 or a subsequence thereof.

9. The peptide of Claim 8 wherein the peptide has an amino acid sequence comprising a subsequence of SEQ ID NO.: 1 and a subsequence of SEQ ID NO.: 2, wherein each subsequence has from five to fifteen amino acid residues.

20 10. The peptide of Claim 8 wherein the peptide has an amino acid sequence consisting of a subsequence of SEQ ID NO.: 1 and a subsequence of SEQ ID NO.: 2, wherein each subsequence has from five to fifteen amino acid residues.

11. The peptide of Claim 10 wherein the subsequence of SEQ ID NO.: 1 and the
25 subsequence of SEQ ID NO.: 2 each have at least nine amino acid residues.


-45-

12. The peptide of Claim 8 wherein the peptide is represented by SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12.
13. A peptide derivative of a peptide represented by SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12, wherein said peptide derivative has angiogenic activity.
14. The peptide derivative of Claim 13 wherein the peptide derivative has between about ten and about twenty-eight amino acid residues or amino acid residue analogs.
15. The peptide derivative of Claim 14 wherein the N-terminus is substituted with an amine protecting group and/or the C-terminus is substituted with a carboxylic acid protecting group.
16. The peptide derivative of Claim 14 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 having from ten to fifteen amino acid residues.
17. The peptide derivative of Claim 14 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 having from about ten to about fifteen amino acid residues, wherein one, two or three amino acid residues in the sequence of the peptide derivative can differ from the amino acid residue(s) in the corresponding position(s) of the subsequence.

-46-

18. The peptide derivative of Claim 14 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, or a subsequence of SEQ ID NO.: 4 or SEQ ID NOS. 9-12, said subsequence having from sixteen to twenty-eight amino acid residues.
- 5 19. The peptide derivative of Claim 14 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3 or a subsequence of SEQ ID NO.: 4 or SEQ ID NOS. 9-12, said subsequence having from about sixteen to about twenty-eight amino acid residues, wherein one four amino, two, three or four acid residues in the sequence of
10 the peptide derivative can differ from the amino acid residue(s) in the corresponding position(s) of the subsequence.
20. The peptide derivative of Claim 14 wherein the peptide derivative is represented by SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12.
21. A multivalent ligand represented by the following structural formula:

15



wherein:

B is a multilinker backbone;

n is an integer from two to about twenty;

each L is a covalent bond or linking group;

-47-

each P is a peptide, wherein at least two of the peptides are a peptide derivative of an AHR of an angiogenic protein, a hybrid peptide, a peptide derivative of a hybrid peptide, or a combination thereof, wherein:

- a) said multivalent ligand has angiogenic activity and
 - 5 b) each peptide and each linker or covalent bond is independently chosen.
22. The multivalent ligand of Claim 21 wherein each L is the same and each peptide is the same.
23. The multivalent ligand of Claim 22 wherein the multilinker backbone is a
10 polymer with two or more reactive groups.
24. The multivalent ligand of Claim 22 wherein the multilinker backbone is a polypeptide.
25. The multivalent ligand of Claim 24 wherein the multilinker backbone is a cascading or pennant polypeptide.
- 15 26. The multivalent ligand of Claim 25 wherein each linker is a covalent bond, a glycine residue or a residue of glycine-glycine.
27. The multivalent ligand of Claim 26 wherein the cascading or pennant polypeptide is a polylysine.
- 20 28. The multivalent ligand of Claim 26 wherein the cascading or pennant polypeptide is a polyaspartic acid or a polyglutamic acid.

-48-

29. The multivalent ligand of Claim 24 wherein the backbone contains a cysteine residue, wherein a bond is formed between the thiol group in the cysteine side chain and a linker or peptide.
30. The multivalent ligand of Claim 27 wherein each peptide is a peptide derivative of the angiogenic homology region of an angiogenic protein.
31. The multivalent ligand of Claim 30 wherein the peptide derivative has between about ten and about twenty-eight amino acid residues.
32. The multivalent ligand of Claim 31 wherein the AHR is represented by SEQ ID NO.: 1 or SEQ ID NO.: 2; and wherein the peptide derivative has between about ten and about twenty-eight amino acid residues or amino acid residue analogs.
33. The multivalent ligand of Claim 32 wherein:
- a) each peptide derivative has between ten and fifteen amino acid residues; and
 - b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2.
34. The multivalent ligand of Claim 32 wherein:
- a) each peptide derivative has between ten and fifteen amino acid residues; and
 - b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2, with the proviso that one, two or three amino acid residues in the

-49-

sequence of the peptide derivative can differ from the amino acid residue(s) in the corresponding position(s) of the subsequence.

35. The multivalent ligand of Claim 32 wherein:
- 5 a) each peptide derivative has between fifteen and twenty-eight amino acid residues; and
- b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2.
36. The multivalent ligand of Claim 32 wherein:
- 10 a) each peptide derivative has between fifteen and twenty-eight amino acid residues; and
- b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2, with the proviso that one, two three or four amino acid residues in the
- 15 sequence of the peptide derivative can differ from the amino acid residue(s) in the corresponding position(s) of the subsequence.
37. The multivalent ligand of Claim 32 wherein:
- a) the multivalent ligand has four peptide derivatives represented by SEQ ID NO.: 1 or SEQ ID NO.: 2; and
- 20 b) the multilinker backbone is a pennant polylysine consisting of three lysine residues.
38. The multivalent ligand of Claim 27 wherein each peptide comprises a sequence of amino acids AA₁ through AA₂₈ or a subsequence thereof having at least ten amino acids, wherein:

-50-

AA₁ is lysine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, glycine or alanine;

5 AA₂ is aspartic acid, asparagine, glutamic acid, glutamine, leucine, valine, isoleucine, methionine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

AA₃ is phenylalanine, alanine, tyrosine, tryptophan or glycine;

AA₄ is threonine, glycine, alanine, cysteine or serine;

AA₅ is alanine, threonine, glycine, cysteine or serine;

10 AA₆ is phenylalanine, tyrosine or tryptophan;

AA₇ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid;

AA₈ tryptophan, alanine, phenylalanine, tyrosine or glycine;

15 AA₉ is arginine, phenylalanine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, lysine, ornithine, tyrosine or tryptophan;

AA₁₀ is leucine, isoleucine, methionine or valine;

AA₁₁ is serine, threonine or alanine;

AA₁₂ is histidine, serine, threonine, cysteine, lysine or ornithine;

20 AA₁₃ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₄ is proline, leucine, valine, isoleucine or methionine;

25 AA₁₅ is lysine, glutamine, histidine, ornithine, asparagine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₆ is threonine, aspartic acid, serine, glutamic acid or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

AA₁₇ is glycine, leucine, alanine, valine, isoleucine or methionine;

-51-

AA₁₈ is phenylalanine, tyrosine or tryptophan;

AA₁₉ is isoleucine, serine, valine, leucine, methionine, cysteine or threonine;

5 AA₂₀ is arginine, isoleucine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline, 2-amino-4-guanidinobutanoic acid, leucine, valine or methionine;

AA₂₁ is methionine, isoleucine, leucine or valine;

10 AA₂₂ valine, arginine, leucine, isoleucine, methionine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₂₃ methionine, arginine, leucine, isoleucine, valine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₂₄ is phenylalanine, alanine, tyrosine, tryptophan or glycine;

15 AA₂₅ is aspartic acid, asparagine, glutamic acid, glutamine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

20 AA₂₆ is glycine, arginine, alanine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₂₇ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid; and

25 AA₂₈ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

wherein each peptide and each linker or covalent bond is independently chosen and the multivalent ligand has angiogenic activity.

-52-


39. The multivalent ligand of Claim 38 wherein the peptide has a sequence comprising a subsequence of SEQ ID NO.: 1 and a subsequence of SEQ ID NO.: 2, wherein each subsequence has from five to fifteen amino acid residues.
- 5 40. The multivalent linker of Claim 39 wherein the peptide has a sequence consisting of a subsequence of SEQ ID NO.: 1 and a subsequence of SEQ ID NO.: 2, wherein each subsequence has from five to fifteen amino acid residues.
- 10 41. The multivalent linker of Claim 40 wherein the subsequence of SEQ ID NO.: 1 and the subsequence of SEQ ID NO.: 2 each have about nine amino acid residues.
- 15 42. The multivalent linker of Claim 38 wherein:
a) the multilinker backbone is a pennant polylysine consisting of three lysine residues;
b) the multivalent ligand comprises four peptides represented by SEQ ID NO. 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12; and
c) each linker is a covalent bond.
43. The multivalent ligand of Claim 27 wherein each peptide is a hybrid peptide or a peptide derivative thereof.
- 20 44. The multivalent ligand of Claim 43 wherein the peptide is represented by SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 or a peptide derivative of SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12.

-53-

45. The multivalent ligand of Claim 44 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 having from ten to fifteen amino acid residues.
- 5 46. The multivalent ligand of Claim 44 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 having from about ten to about fifteen amino acid residues, wherein one, two or three amino acid residue(s) in the sequence of the peptide derivative can differ from the amino acid residue(s) in the corresponding position(s) of the subsequence.
- 10
47. The multivalent ligand of Claim 44 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3 or a subsequence of SEQ ID NO.: 4 or SEQ ID NOS. 9-12, said subsequence having from sixteen to twenty-eight amino acid residues.
- 15 48. The multivalent ligand of Claim 44 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, a subsequence of SEQ ID NO.: 4, or SEQ ID NOS. 9-12 said subsequence having from about sixteen to about twenty-eight amino acid residues, wherein one, two, three or four amino acid residue(s) in the sequence of the peptide derivative can differ from the amino acid residue(s) in the corresponding position(s) of the subsequence.
- 20
49. A method of modulating angiogenesis in a subject comprising administering a therapeutically effective amount of a multivalent ligand having angiogenic

-54-

activity, said multivalent ligand being represented by the following structural formula:



wherein:

- 5 B is a multilinker backbone;
 n is an integer from two to about twenty;
 each L is a covalent bond or linking group;
 each P is a peptide, wherein at least two of the peptides are a peptide
 derivative of an AHR of an angiogenic protein, a hybrid peptide or a peptide
 10 derivative of a hybrid peptide, or a combination thereof, wherein:
 a) said peptide has angiogenic activity; and
 b) each peptide and each linker or covalent bond is independently
 chosen.
50. The method of Claim 49 wherein the multivalent ligand is anti-angiogenic
 15 and the subject is being treated for cancer.
51. The method of Claim 49 wherein the multivalent ligand is pro-angiogenic.
52. The method of Claim 49 wherein:
 a) the multivalent peptide backbone is a pennant or cascading
 polylysine;
 20 b) each linker and each polypeptide is the same;

-55-

- c) and each linker is a covalent bond, a glycine residue or a glycine-glycine residue.
53. The method of Claim 52 wherein each peptide is a peptide derivative of the AHR of an angiogenic protein.
- 5 54. The method of Claim 53 wherein:
- a) the angiogenic homology region is represented by SEQ ID NO.: 1 or SEQ ID NO.: 2; and
 - b) the peptide derivative has between about ten and about twenty-eight amino acid residues or amino acid residue analogs.
- 10 55. The method of Claim 54 wherein:
- a) each peptide derivative has between ten and fifteen amino acid residues; and
 - b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2.
- 15 56. The method of Claim 54 wherein:
- a) each peptide derivative has between ten and fifteen amino acid residues; and
 - b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2,
- 20 with the proviso that one, two or three amino acid residues in the sequence of the peptide derivative can differ from the amino acid residue(s) in the corresponding position(s) in the subsequence.

-56-

57. The method of Claim 55 wherein:
- a) each peptide derivative has between fifteen and twenty-eight amino acid residues; and
 - b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2.
58. The method of Claim 54 wherein:
- a) each peptide derivative has between fifteen and twenty-eight amino acid residues; and
 - b) each peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 1 or a subsequence of SEQ ID NO.: 2, with the proviso that one, two, three or four amino acid residues in the sequence of the peptide derivative can differ from the amino acid residue(s) in the corresponding position of the subsequence.
59. The method of Claim 54 wherein the multilinker backbone is a pennant polylysine consisting of four amino acids, and the peptide derivative is represented by SEQ ID NO.: 1 or SEQ ID NO.: 2.
60. The method of Claim 52 wherein the peptide comprises a sequence of amino acids AA₁ through AA₂₈ or a subsequence thereof having at least ten amino acids, wherein:
- AA₁ is lysine, ornithine, arginine, N-nitroarginine, β-cycloarginine, γ-hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, glycine or alanine;
- AA₂ is aspartic acid, asparagine, glutamic acid, glutamine, leucine, valine, isoleucine, methionine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

-57-

AA₃ is phenylalanine, alanine, tyrosine, tryptophan, leucine, isoleucine, methionine, valine or glycine;

AA₄ is threonine, glycine, alanine, cysteine or serine;

AA₅ is alanine, threonine, glycine, cysteine or serine;

5 AA₆ is phenylalanine, tyrosine or tryptophan;

AA₇ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid;

AA₈ tryptophan, alanine, phenylalanine, tyrosine or glycine;

10 AA₉ is arginine, phenylalanine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline 2-amino-4-guanidinobutanoic acid, lysine, ornithine, tyrosine or tryptophan;

AA₁₀ is leucine, isoleucine, methionine or valine;

AA₁₁ is serine, threonine or alanine;

AA₁₂ is histidine, serine, threonine, cysteine, lysine or ornithine;

15 AA₁₃ is arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₄ is proline, leucine, valine, isoleucine or methionine;

20 AA₁₅ is lysine, glutamine, histidine, ornithine, asparagine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₁₆ is threonine, aspartic acid, serine, glutamic acid or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

AA₁₇ is glycine, leucine, alanine, valine, isoleucine or methionine;

AA₁₈ is phenylalanine, tyrosine or tryptophan;

25 AA₁₉ is isoleucine, serine, valine, leucine, methionine, cysteine or threonine;

-58-

AA₂₀ is arginine, isoleucine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline, 2-amino-4-guanidinobutanoic acid, leucine, valine or methionine;

AA₂₁ is methionine, isoleucine, leucine or valine;

5 AA₂₂ valine, arginine, leucine, isoleucine, methionine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

10 AA₂₃ methionine, arginine, leucine, isoleucine, valine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

AA₂₄ is phenylalanine, alanine, tyrosine, tryptophan or glycine;

AA₂₅ is aspartic acid, asparagine, glutamic acid, glutamine or a substituted or unsubstituted aliphatic or aryl ester of glutamic acid or aspartic acid;

15 AA₂₆ is glycine, arginine, alanine, ornithine, lysine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

20 AA₂₇ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid; and

AA₂₈ is lysine, alanine, arginine, glycine, serine, ornithine, arginine, N-nitroarginine, β -cycloarginine, γ -hydroxyarginine, N-amidinocitruline or 2-amino-4-guanidinobutanoic acid;

25 wherein each peptide and each linker or covalent bond is independently chosen and the multivalent ligand has angiogenic activity.

-59-


61. The method of Claim 60 wherein the peptide has a sequence comprising a subsequence of SEQ ID NO.: 1 and a subsequence of SEQ ID NO.: 2, wherein each subsequence has from five to fifteen amino acid residues.
- 5 62. The method of Claim 61 wherein the peptide has a sequence consisting of a subsequence of SEQ ID NO.: 1 and a subsequence of SEQ ID NO.: 2, wherein each subsequence has from five to fifteen amino acid residues.
63. The method of Claim 62 the subsequence of SEQ ID NO.: 1 and the subsequence of SEQ ID NO.: 2 each have at least nine amino acid residues.
- 10 64. The method of Claim 52 wherein the peptide is a hybrid peptide or a peptide derivative thereof.
65. The method of Claim 64 wherein the hybrid peptide is represented by SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 or is a peptide derivative thereof.
- 15 66. The method of Claim 65 wherein the peptide derivative of SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 have between about ten and about twenty-eight amino acid residues or amino acid residue analogs.
67. The method of Claim 66 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 having from ten to fifteen amino acid residues.
- 20 68. The method of Claim 66 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, SEQ ID NO.: 4

-60-

or SEQ ID NOS. 9-12 having from about ten to about fifteen amino acid residues, wherein one, two or three amino acid residues in the sequence of the peptide derivative can differ from the amino acid residue(s) at the corresponding positions(s) of the subsequence.

- 5 69. The method of Claim 66 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3 or a subsequence of SEQ ID NO.: 4 or SEQ ID NOS. 9-12, said subsequence having from sixteen to twenty-eight amino acid residues.
- 10 70. The method of Claim 66 wherein the peptide derivative has an amino acid sequence corresponding to a subsequence of SEQ ID NO.: 3, a subsequence of SEQ ID NO.: 4 or SEQ ID NOS. 9-12, said subsequence having from about sixteen to about twenty-eight amino acid residues, wherein one, two, three or four amino acid residues in the sequence of the peptide derivative can differ from the amino acid residue(s) at the corresponding positions(s) of the subsequence.
- 15 71. The method of Claim 66 wherein the peptide derivative is represented by SEQ ID NO.: 3, SEQ ID NO.: 4 or SEQ ID NOS. 9-12 and wherein the multilinker backbone is a pennant polylysine consisting of three lysine residues.
- 20 72. A polypeptide multivalent ligand represented by the following structural formula:

-61-



wherein:


m is an integer from zero to about twenty;

each S is a peptide spacer having from about five to about thirty
5 amino acids;

each P is a peptide having from about ten to about forty amino acid
residues, wherein at least two of the peptides are peptide derivatives of an
AHR of an angiogenic protein, a hybrid peptide or a peptide derivative of a
hybrid peptide, wherein:

- 10 a) said polypeptide multivalent ligand has angiogenic activity; and
b) each peptide and each peptide spacer is independently chosen.

73. A method of modulating angiogenesis in a subject comprising administering
a therapeutically effective amount of a polypeptide multivalent ligand having
15 angiogenic activity, said polypeptide multivalent ligand being represented by
the following structural formula:



wherein:

-62-

m is an integer from zero to about twenty;

each S is a peptide spacer having from about five to about thirty amino acids;

5 each P is a peptide having from about ten to about thirty amino acid residues, wherein at least two of the peptides are peptide derivatives of an AHR of an angiogenic protein, a hybrid peptide or a peptide derivative of a hybrid peptide, wherein:

- a) said polypeptide multivalent ligand has angiogenic activity; and
- b) each peptide and each peptide spacer is independently chosen.

TSP-1	R H I G W K D F T A Y R R L S H R P K T G F I R V V M Y E G K K I M	-5	1	5	10	15	20	25
Endostatin	Q A R A V G L A G T F R A F L S S R L Q D L Y S I V R R A D R A A V W							
TSP-4	R N V G W K D K V S Y R W F L Q H R P Q V G Y I R V R F Y E G S E L V							
Angiostatin	I S K T M S G L E C Q A W D S Q S P H A H G Y I P S K F P N K N I K K							

Fig. 1

SEQ ID NO. 3	A C - T F R A F L S S R L Q T G F I R V V M Y E G	1	5	10	15	20
SEQ ID NO. 4	A C - A Y R W R L S H R P K D L Y S I V R R A D G					

Fig. 2

SEQ ID NO. 7	1	5	10	15	20
	AC-T	F R A F L S S R L	Q D L Y S I	V R R A D G	
SEQ ID NO. 8	AC-A	Y R W R L S H R P	K T G F I R V	V M Y E G	
SEQ ID NO. 9	AC-T	A Y R W R L S H R P	K D L Y S I	V R R A D R	
SEQ ID NO. 10	AC-A	Y R W R L S H R P	K D L Y S I	V R R A D R	
SEQ ID NO. 11		AC-R	W R L S H R P	K D L Y S I	V R R A D R
SEQ ID NO. 12	AC-K-D-F-T-A	Y R W R L S H R P	K D L Y S I	V R R A D R	

Fig. 3

R/Q	H/A	I/R	G/A	V/W	K/G	D/L	F/A	T/G	A/T	Y/F	R	W/A	R/F	L
				-5					1				5	
		S	S/H	R	P/L	K/Q	T/D	G/L	F/Y	I/S	R/I			
					10					15				
V	R/V	M/R	Y/A	E/D	G/R	K/A	I/V	M/W	A/I					
			20					25						

Fig. 5




Fig. 4A




Fig. 4B




Fig. 4C





Fig. 4D






FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11





Fig. 12

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/06246

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K14/78 C12N9/68 C07K14/515 A61K47/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 399 667 A (FRAZIER & KOSFELD) 21 March 1995 (1995-03-21) figure 2A	9
X	WO 95 05191 A (UAB RESEARCH FOUNDATION) 23 February 1995 (1995-02-23) claim 4	9
X	WO 96 17059 A (WASHINGTON UNIVERSITY) 6 June 1996 (1996-06-06) figure 1	9
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 July 1999

Date of mailing of the international search report

29/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Masturzo, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/06246

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>D J GOOD ET AL.: "A tumor suppressor-dependent inhibitor of angiogenesis is immunologically and functionally indistinguishable from a fragment of thrombospondin "</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA.,</p> <p>vol. 87, September 1990 (1990-09), pages 6624-6628, XP002109500</p> <p>NATIONAL ACADEMY OF SCIENCE. WASHINGTON., US</p> <p>ISSN: 0027-8424</p> <p>table 1</p> <p>-----</p>	1-74

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/06246

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5399667	A	21-03-1995	US 5627265 A	06-05-1997
WO 9505191	A	23-02-1995	AU 697624 B	15-10-1998
			AU 7565294 A	14-03-1995
			CA 2169360 A	23-02-1995
			EP 0716609 A	19-06-1996
			JP 9505555 T	03-06-1997
WO 9617059	A	06-06-1996	US 5627265 A	06-05-1997
			AU 4151496 A	19-06-1996
			CA 2206553 A	06-06-1996
			EP 0795013 A	17-09-1997
			JP 10509974 T	29-09-1998